

Immunization Timing 2024



Birth

HepB¹

RSV²

(age: 0-8 months)

**Age
2
months**

Interval
from
previous
dose

DTaP

(Diphtheria,
Tetanus,
Pertussis)

**Polio
(IPV)**

HepB³

(age: 1-2
months)

1-2
months
after
birth dose

Hib

(Hib meningitis)

PCV

(Pneumo)

RV⁴

(Rotavirus)

**Age
4
months**

Interval
from
previous
dose

DTaP

1-2
months

**Polio
(IPV)**

1-2
months

HepB³

if 1st dose
given at
2 months

1-2
months

Hib

1-2
months

PCV

1-2
months

RV⁴

4-10
weeks

**Age
6
months**

Interval
from
previous
dose

DTaP

1-2
months

Polio
age: 6-18
months)

1-14
months

HepB³
(age: 6-18
months)

2-12
months
and
≥4 months
after
1st dose

Hib⁵

1-2
months

PCV

1-2
months

RV⁴

if RotaTeq
used for
doses 1 or 2

4-10
weeks

**Age
12
months**

Interval
from
previous
dose

HepA⁸
(age: 12-23
months)

MMR^{8,9,10}
(ages 12-15
months)

Var¹⁰

(age: 12-15
months)

Hib

(age: 12-15
months)

PCV¹¹

(age: 12-15
months)

Interval
from
previous
dose

**Age
15
months**

Interval
from
previous
dose

DTaP¹²

6-12
months

**Age
18
months**

Interval
from
previous
dose

HepA

6-18
months

**Age
4-6
years**

DTaP

Polio (IPV)

MMR^{9, 10}

Varicella¹⁰

**Age
11-12
years**

HPV¹³ (2 doses, can
start at age 9)

MenACWY (MCV4)

Tdap

**Age
16
years**

MenACWY (MCV4)

MenB¹⁴

6 months – 18+ years

COVID-19 vaccine(s)⁶

Flu vaccine, every fall⁷